

Centro-symmetric Hamiltonians foster quantum transport

Mattia Walschaers,^{1,2,*} Roberto Mulet,^{1,3,†} and Andreas Buchleitner^{1,‡}

¹*Physikalisches Institut, Albert-Ludwigs-Universität Freiburg,
Hermann-Herder-Str. 3, D-79104 Freiburg, Germany*

²*Instituut voor Theoretische Fysica, KU Leuven,
Celestijnenlaan 200D, B-3001 Heverlee, Belgium*

³*Complex System Group, Department of Theoretical Physics, University of Havana, Cuba*

(Dated: July 18, 2012)

We propose a model for fast and highly efficient quantum transport of excitations, through finite, disordered systems. The presented mechanism is statistically robust against configurational changes which alter the realization of disorder. We finally discuss the potential relevance of our findings for excitation transport in photosynthetic light harvesting complexes.

PACS numbers: 05.60.Gg, 03.65.Xp, 72.10.-d, 87.15.hj

In a variety of fields, ranging from quantum information [1] to solar cell physics [2], the efficient transport of quanta is of paramount importance. Under which general conditions fundamental principles of quantum mechanics can be exploited to enhance transport in complex systems remains a widely open question. Common wisdom suggests that quantum interference can enhance transport across perfectly periodic potentials [3, 4], while it tends to suppress transport in disordered systems [5, 6]. In general, multi-path quantum interference leads to erratic fluctuations of transmission probabilities when boundary conditions are slightly changed [7–11]. These fluctuations are often indicative of the strong, non-linear coupling of few degrees of freedom, as it abounds in heavy nuclei [7], ultra-cold many-particle dynamics [12], strongly perturbed Rydberg systems [13–15] billiard geometries for photons [16] and electrons [17], strongly driven quantum systems [18], and in large molecules [19, 20]. Can we identify structural properties of the underlying Hamiltonians in those specific cases which exhibit *quantum-enhanced* transport? Are these properties statistically robust, in the sense that they are “implementable” by controlling only few coarse-grained parameters, without claiming full control of the detailed structure of a possibly large, composite quantum system? It is the present contribution’s purpose to offer an affirmative answer to these questions, on the basis of very general considerations. As a possible application, we will discuss the potential role of our findings for efficient light harvesting in photosynthetic complexes.

Let us start with the definition of our model, which is inspired by the structural elements of the light harvesting complex FMO of sulfur bacteria [21, 22] and by the theory of quantum graphs [23]: We consider the coherent quantum transport of one single excitation across a disordered network of N sites. Hilbert space is spanned by the basis states $|i\rangle$ which represent those states where the excitation is fully localized at the network’s site i . In order to formulate a quantitative, statistical theory, we generate different realizations of disorder by sampling

over $N \times N$ Hamiltonians H from the Gaussian Orthogonal Ensemble [24], with matrix entries $H_{i,j}$ which encode the couplings between sites i and j . For each realization, input $|\text{in}\rangle$ and output $|\text{out}\rangle$ are defined as those sites with the weakest coupling $V = \min_{i \neq j} |H_{i,j}|$. Our figure of merit is the transfer efficiency

$$\mathcal{P}_H = \max_{t \in [0, T_R]} |\langle \text{out}, \phi(t) \rangle|^2, \quad |\phi(0)\rangle = |\text{in}\rangle, \quad (1)$$

which quantifies a given random network’s performance in terms of excitation transport from $|\text{in}\rangle$ to $|\text{out}\rangle$. $\mathcal{P}_H$ is gauged against the direct coupling V between $|\text{in}\rangle$ to $|\text{out}\rangle$, in the absence of all intermediate sites, through the definition of the associated benchmark time scale $T_R = \pi/2V$ [22, 25].

We have shown earlier, in a study of 3D random graphs with dipole-dipole coupling [26] that a strong and unambiguous correlation exists between high transfer efficiencies and the Hamiltonian’s vicinity in matrix space to a *centro-symmetric* structure with respect to $|\text{in}\rangle$ and $|\text{out}\rangle$, which is defined by $JH = HJ$ and $|\text{in}\rangle = J|\text{out}\rangle$ [27]. J is the exchange matrix, $J_{i,j} = \delta_{i, N-j+1}$ [28]. However, we did also observe that even for high centro-symmetries the distribution of transfer efficiencies was still rather broad, implying that centro-symmetry alone is not sufficient for efficient state transfer. This can be easily understood by the fact that, while centro-symmetry does distinguish $|\text{in}\rangle$ and $|\text{out}\rangle$ from the intermediate sites, it does not exclude network conformations which trap the excitation in the vicinity of $|\text{in}\rangle$, and thus inhibit transmission. Therefore, we need to identify an additional structural element which guarantees robustness, in the sense that the transfer efficiency must not depend strongly on the specific conformation of the intermediate sites. Note that such a feature is also of obvious relevance for our model to be applicable to realistic light harvesting complexes, which indeed do undergo structural changes on the macromolecular scale.

Intuitively, this structural stability of efficient excitation transfer from $|\text{in}\rangle$ to $|\text{out}\rangle$ is guaranteed if both states

are coupled through a dominant tunneling doublet in the spectrum. The sole role of the intermediate states is then to *collectively amend* the effective tunneling coupling by a collective energy shift Δs . If Δs has the proper sign, this can lead to a dramatic enhancement of the transfer efficiency. Note that such collective shifts induced by the coupling to random or “chaotic” states have been investigated in the context of chaos assisted tunneling (CAT) [29–31]. The existence of a dominant doublet will enter as a key hypothesis into the subsequent analytical description of our problem.

Given the centro-symmetry of H , it can be cast, through an orthogonal transformation to the eigenbasis of the exchange operator J , into the block diagonal representation [28]

$$H = \begin{pmatrix} H^+ & 0 \\ 0 & H^- \end{pmatrix}. \quad (2)$$

In this new form, both H^+ and H^- are again $N/2 \times N/2$ GOE matrices, *i.e.* the elements $H_{i,j}^\pm$ are sampled from a Gaussian distribution with zero mean and variance $(1 + \delta_{i,j})2\xi^2/N$.

Since two of the eigenvectors of J have the form

$$|\pm\rangle = \frac{1}{\sqrt{2}}(|\text{in}\rangle \pm |\text{out}\rangle), \quad (3)$$

we now additionally assume (see above) that $|+\rangle$ and $|-\rangle$ form a dominant doublet, such that they are both close to eigenstates $|\tilde{+}\rangle$ and $|\tilde{-}\rangle$ of H^+ and H^- , respectively. It is then useful to write the Hamiltonian (2) as

$$H = \begin{pmatrix} E+V & \langle \mathcal{V}^+ | & & \\ | \mathcal{V}^+ \rangle & H_{sub}^+ & & \\ & & E-V & \langle \mathcal{V}^- | \\ | \mathcal{V}^- \rangle & & & H_{sub}^- \end{pmatrix}, \quad (4)$$

which makes the definition of rows and columns which relate to $|+\rangle$ and $|-\rangle$ explicit. From the definition of $|\pm\rangle$ it is easy to see that $\langle H \rangle_\pm = E \pm V$. $|\mathcal{V}^\pm\rangle$ contains the couplings of the dominant doublet states $|\pm\rangle$ to the remainder of the system.

Since the norm $\|\mathcal{V}^\pm\|$ of the coupling is small, due to the assumption of the dominant doublet,

$$|\langle \tilde{\pm}, \pm \rangle|^2 > \alpha \approx 1. \quad (5)$$

Under this condition, perturbation theory guarantees that $E \pm V$ in (4) are eigenvalues of H , up to some small correction term $s^\pm$. The explicit expression for the transfer efficiency is then dominated by those terms associated with $|\tilde{\pm}\rangle$, leading to the estimate

$$\mathcal{P}_H \gtrsim \max_{t \in [0, T_R]} \frac{\alpha^2}{4} \left| e^{-it(E+V+s^+)} - e^{-it(E-V+s^-)} \right|^2, \quad (6)$$

where $s^\pm = \sum_i \frac{|\langle \mathcal{V}^\pm, \psi_i^\pm \rangle|^2}{E \pm V - e_i^\pm},$

and $|\psi_i^\pm\rangle$ and $e_i^\pm$ are the eigenvectors and eigenvalues of $H_{sub}^\pm$. From (6) it is clear that the efficiency is large, $\mathcal{P}_H \gtrsim \alpha^2$, if $t = \pi/|2V + \Delta s|$ is smaller than T_R , and we can interpret $|2V + \Delta s|$ as an effective tunneling rate. All realizations for which we obtain $T_R/t > 1$ have efficient transport which is faster than the direct coupling between $|\text{in}\rangle$ and $|\text{out}\rangle$.

For fixed E and V the distribution of Δs was already derived within the context of the CAT [30, 31], and it turned out that for large N it is a Cauchy (or Lorentzian) distribution. In our present problem, E and V are stochastic variables, and we should average the resulting distribution over E and V . Considering that the integrations over E and V are dominated by the mean values of their distributions, which are given by $\overline{V} \approx \sqrt{2\pi\xi}/N^{3/2}$ and $\overline{E^2} = 2\xi^2/N$ (since $\overline{E} = 0$, we use $\overline{E^2}$), a lengthy but straightforward calculation shows that the probability distribution of $T_R(2V + \Delta s)/\pi = T_R/t$ is given by

$$P\left(\frac{T_R}{t} = x\right) = \frac{1}{\pi} \left(\frac{s_0}{s_0^2 + (1+x)^2} + \frac{s_0}{s_0^2 + (1-x)^2} \right),$$

with $s_0 = \frac{\overline{\|\mathcal{V}\|^2} N^{1/2} (2 - 1/2N)}{\sqrt{2\pi\xi^2}}.$ (7)

Here, $\overline{\|\mathcal{V}\|^2}$ denotes the expectation value of $\|\mathcal{V}^\pm\|^2$ for all realizations where the dominant doublet assumption holds.

Under the prerogative of a dominant doublet, the distribution (7) depends on only two parameters: ξ characterizes the spectral density of the eigenstates of H_{sub}^+ and H_{sub}^- , while $\overline{\|\mathcal{V}\|^2}$ measures the average coupling strength of the dominant doublet to these states. It therefore cannot be emphasized enough that, within the picture here elaborated, the transport properties of the problem *do not* depend on the specificities of the intermediate electronic states of the network – *i.e.* on its specific structural conformation.

In order to validate our theoretical model by numerical simulations, we generate many GOE Hamiltonians with the additional constraint of centro-symmetry with respect to $|\text{in}\rangle$ and $|\text{out}\rangle$. For each of these Hamiltonians the existence of a dominant doublet is assessed by inspection of its eigenvectors and by verifying that, for some $|\tilde{\pm}\rangle$, the condition $|\langle \tilde{\pm}, \pm \rangle|^2 > \alpha$ holds. Then, the quantity $\mathcal{P}_H$ is obtained by numerically solving the quantum dynamics of the Hamiltonian, and t is defined as the earliest point in time for which $|\langle \text{out}, \phi(t) \rangle|^2 = \mathcal{P}_H$.

Figure 1 compares numerical results and analytical prediction (7). It must be noticed that there are no free parameters involved in this plot: The values $\xi = 2$, $\alpha > 0.95$ and $\overline{\|\mathcal{V}\|^2} = 0.323192$ are either given a priori or directly extracted from the statistical characterization of the numerically generated sample Hamiltonians. Clearly, the majority of realizations have t smaller than




FIG. 1: Histogram of the numerically simulated inverse transfer time T_R/t (dashed line), compared to the theoretical prediction (7) (full line), for density of states $\xi = 2$, $N = 10$ network sites, and a dominant doublet condition $\alpha = 0.95$ (see (5)). $\|\mathcal{V}\|^2 = 0.323192$ is directly extracted from the numerical sampling over different random Hamiltonians, and the evident agreement between simulation and theory results with no adjustable parameters. Simulations were performed in the time window $[0, 1.7T_R]$; therefore the plot only starts at $T_R/t = (1.7)^{-1}$. The distribution's heavy tail (as highlighted in the inset, on a log scale) guarantees the abundance of very short transfer times $t \ll T_R$ as compared to the direct coupling time between $|\text{in}\rangle$ and $|\text{out}\rangle$.

the time scale set by the direct coupling V between $|\text{in}\rangle$ and $|\text{out}\rangle$. The fat algebraic tail of the Cauchy distribution for $t \ll T_R$ guarantees that realizations with very fast transport, with respect to T_R , are abundant in the sense that they are not exponentially unlikely.

We still need to verify that the such defined Hamiltonians provide efficient transport. For a fixed $\alpha = 0.95$, our model predicts that the efficiency should be larger than $\mathcal{P}_H \gtrsim \alpha^2 \approx 0.9$. This is indeed observed in the simulations. Figure 2 shows the probability distribution of the efficiencies, which is sharply peaked above $\mathcal{P}_H > 0.9$. Comparison, in the same figure, with the probability distributions of the efficiencies of centro-symmetric GOE matrices without the doublet constraint, and of general GOE matrices, respectively, shows that in both cases the average efficiency is significantly lower than for these centro-symmetric Hamiltonians which exhibit a dominant doublet.

A remarkable asset of the transport mechanism here presented is its robustness under different realizations of disorder, which, in the context of networks, refers to different configurations of the intermediate sites (represented by $H_{\text{sub}}^\pm$ in (4)). In the light of the recent debate on the potential role of quantum coherence in photosynthetic harvesting of the sunlight's energy, one may wonder whether this mechanism is already implemented by Nature. We now briefly scrutinize such hypothesis, on




FIG. 2: Histograms of the transfer efficiencies $\mathcal{P}_H$ for three different Hamiltonian matrix ensembles, with fixed density of states $\xi = 2$ and number $N = 10$ of network sites. Clearly, as we progressively constrain the ensemble from GOE to centro-symmetric GOE and, finally, to centro-symmetric with dominant doublet condition $\alpha = 0.95$, the average efficiencies are dramatically enhanced. In perfect accord with our analytical model, the efficiency distribution is strongly peaked at values $\mathcal{P}_H \geq \alpha^2 \approx 0.9$ (indicated by the arrow), when we only sample over random Hamiltonians with dominant doublet.

the basis of published data on the widely studied FMO light harvesting complex.

To start with, the recently discovered eighth site of the FMO molecular network is only weakly coupled to the remainder of the complex [21]. This is indeed consistent with the above intuition of a dominant doublet. Furthermore, the centro-symmetry of the published FMO Hamiltonian H_0 [21] was studied in [26], and it was found that, while H_0 is itself neither efficient nor centro-symmetric, weak perturbations of the matrix elements of H_0 allow for high efficiencies [32], together with enhanced centro-symmetries [26]. Consistently, verification of the dominant doublet condition (5) for H_0 [33] results in $\max_i |\langle \pm, \eta_i \rangle|^2 \approx 0.5$, upon maximization over all eigenvectors $|\eta_i\rangle$. In contrast, the optimized Hamiltonian inferred in [26] indeed does exhibit a dominant doublet, with $\max_i |\langle +, \eta_i \rangle|^2 = 0.877361$ and $\max_i |\langle -, \eta_i \rangle|^2 = 0.901908$, respectively.

In conclusion, we described a general mechanism that gives rise to fast and efficient quantum transport in finite, disordered systems. The mechanism rests on two crucial ingredients: The first is the centro-symmetry of the underlying Hamiltonian, which guarantees the existence of a block diagonal representation. The second is the existence of a dominant doublet, that guarantees the statistical robustness of the transport properties, under the coupling to random/chaotic states which assist the transport. Note, in particular, that the structure of (4) is maintained under averages over different realizations

of centro-symmetric matrices with dominant doublets (what provides an explanation for the on a first glance surprising numerical observation in [26], that also averages over efficient Hamiltonians remain efficient). Both these features appear consistent with published effective Hamiltonians of the FMO light harvesting complex and weak perturbations thereof. The statistics of the transfer efficiencies and times as shown in Figs. 1 and 2 only depend on the intermediate network sites' density of states ξ , and on the average coupling strength $\|\overline{\mathcal{V}}\|^2$ of the in- and output-sites to the network. On the one hand, this may explain why coherence effects can survive even in relatively floppy macromolecular structures, if only these *average* quantities can be stabilized by the functional units' architecture. On the other hand, if such stabilization is possible, this immediately implies a handle of control, by proper tuning of ξ , $\|\overline{\mathcal{V}}\|^2$ and, possibly, N [21], according to the physiological needs.

We enjoyed stimulating discussions with Mark Fannes, Thomas Wellens, Yoshitaka Tanimura, Maciej Zworski, Anthony Dutoi, and Rienk van Grondelle. R.M. is thankful to the Alexander von Humboldt Stiftung for support through a Research Fellowship for Experienced Researchers. M.W. and A.B. are grateful for DFG funding under the project "Scattering systems with complex dynamics". M.W. acknowledges partial funding by Belgian Interuniversity Attraction Poles Programme P6/02 and FWO Vlaanderen project G040710N.

* Electronic address: mattia@itf.fys.kuleuven.be

† Electronic address: roberto.mulet@gmail.com

‡ Electronic address: abu@uni-freiburg.de

- [1] M. Christandl, N. Datta, A. Ekert and A. J. Landahl, Phys. Rev. Lett. **92**, 187902 (2004).
- [2] J. Nelson, *The Physics of Solar Cells*, Imperial College Press, London, 2010.
- [3] N. W. Ashcroft and N. D. Mermin, *Solids State Physics*, Saunders College Publishing, Philadelphia, 1976.
- [4] T. Aharonov, L. Davidovich, and N. Zagury, Phys. Rev. A **48**, 1687 (1993).
- [5] P. W. Anderson, Phys. Rev. **109**, 14921505 (1958).
- [6] G. Modugno, Rep. Prog. Phys. **73**, 102401 (2010).
- [7] G. E. Mitchell, A. Richter, H. A. Weidenmüller, Rev. Mod. Phys. **82**, 2845 (2010).
- [8] B. Kramer and A. MacKinnon, Rep. Prog. Phys. **56**, 1469 (1993).
- [9] J. Madroñero, A. Ponomarev, A.R.R. Carvalho, S. Wimberger, C. Viviescas, A. Kolovsky, K. Hornberger, P. Schlagheck, A. Krug, and A. Buchleitner, Adv. At. Mol. Opt. Phys. **53**, 33 (2006).
- [10] T. Dittrich, P. Hänggi, G.L. Ingold, B. Kramer, G. Schön, W. Zwerger (eds.), *Quantum Transport and Dissipation*, Wiley-VCH, Weinheim, 1998.
- [11] S. Löck, A. Bäcker, R. Ketzmerick, and P. Schlagheck, Phys. Rev. Lett. **104**, 114101 (2010).
- [12] S. Hunn, M. Hiller, D. Cohen, T. Kottos and A. Buchleitner, J. Phys. B **45**, 085302 (2012).
- [13] N.N. Choi, M.H. Lee, and G. Tanner, Phys. Rev. Lett. **93**, 054302 (2004).
- [14] J. Madroñero and A. Buchleitner, Phys. Rev. Lett. **95**, 263601 (2005).
- [15] J.H. Gurian, P. Cheinet, P. Huillery, A. Fioretti, J. Zhao, P.L. Gould, D. Comparat, P. Pillet, Phys. Rev. Lett. **108**, 023005 (2012).
- [16] H.-J. Stöckmann and J. Stein, Phys. Rev. Lett. **64**, 2215 (1990).
- [17] A. Sachrajda, R. Ketzmerick, C. Gould, Y. Feng, P. Kelly, A. Delage, Z. Wasilewski, Phys. Rev. Lett. **80**, 001948 (1998).
- [18] A.R.R. Carvalho and A. Buchleitner, Phys. Rev. Lett. **93**, 204101 (2004).
- [19] S. Keshavamurthy and P. Schlagheck, *Dynamical Tunneling - Theory and Experiment*, CRC Press, Boca Raton, 2011.
- [20] T.P.J. Krüger, E. Wientjesb, R. Crocea, and R. v. Grondelle, Proc. Nat. Acad. Sci. **108**, 13516 (2011).
- [21] M. Schmidt am Busch, F. Müh, M. E. Madjet and T. Renger, J. Phys. Chem. Lett. **2**, 93 (2011).
- [22] T. Scholak, F. Mintert, T. Wellens, and A. Buchleitner, Semicond. Semimet. **83**, 1 (2010).
- [23] T. Kottos and U. Smilansky, Ann. Phys. **274**, 76 (1999).
- [24] M. L. Mehta, *Random Matrices*, Elsevier/Academic Press, Amsterdam, 2004.
- [25] T. Scholak, T. Wellens and A. Buchleitner, Europhys. Lett. **96**, 10001 (2011).
- [26] T. Zech, R. Mulet, T. Wellens and A. Buchleitner, arXiv:1205.5519.
- [27] This constraint actually implies $V = \min_i |H_{i,N-i+1}|$.
- [28] A. Cantoni and P. Butler, Linear Alg. Appl. **13**, 275 (1976).
- [29] S. Tomsovic and D. Ullmo, Phys. Rev. E **50**, 145162 (1994).
- [30] F. Leyvraz and D. Ullmo, J. Phys. A **29**, 2529 (1996).
- [31] J. Zakrzewski, D. Delande and A. Buchleitner, Phys. Rev. E **57**, 1458 (1998).
- [32] T. Scholak, F. de Melo, T. Wellens, F. Mintert, and A. Buchleitner, Phys. Rev. E **83**, 021912 (2011).
- [33] Following the approach of [26], we set all diagonal matrix elements of H_0 to zero, in this analysis.